

Fixed Trammel Nets Interactions with Bottlenose Dolphin

1. Introduction

The Assessing Welsh Fishing Activities (AWFA) Project is a structured risk-based approach to determining impacts from current and potential fishing activities (those undertaken from licensed and registered commercial fishing vessels), upon the features of European marine sites (EMS) in Wales.

Further details of the AWFA project, and all completed assessments to date, can be found on the [AWFA website](#).

The methods and process used to classify the risk of interactions between fishing gears and EMS features, as purple (high), orange (medium) or green (low) risk, can be found in the AWFA Project Phase 1 outputs: [Principles and Prioritisation Report](#) and resulting [Matrix](#) spreadsheet.

2. Assessment summary

<p>Assessment Summary:</p> <p>Fixed Trammel Nets</p> <p>Interactions with</p> <p>Bottlenose Dolphin</p>	<p>No studies were found that directly measured or estimated impacts of fixed trammel nets on bottlenose dolphin, however indirect evidence from other fixed net fisheries were considered.</p> <p><u>Assessment of impact pathway 1. Direct capture, damage, disturbance or harm to a designated species feature:</u></p> <p>The impacts from fixed nets or noise pollution associated with fishing vessels could lead to bottlenose dolphin bycatch, displacement or disturbance.</p> <p><u>Assessment of impact pathway 2. Damage to the habitat of designated species features (including through direct physical impact, pollution, changes in thermal regime, hydrodynamics, light etc.):</u></p> <p>The impacts from nets, weights or anchors are not likely to affect the integrity of the water column habitats utilised by bottlenose dolphin (see Impact Pathway 4 for consideration of benthic feeding activity and prey habitat).</p> <p><u>Assessment of impact pathway 3. Removal of prey species of a designated species feature:</u></p>
---	--

	<p>The removal of prey species by fixed trammel nets could affect bottlenose dolphins. However, evidence suggests that bottlenose dolphins will readily switch prey, but it is not known if dependency on alternative prey availability and quality is detrimental at the population level in the long term</p> <p><u>Assessment of impact pathway 4. Damage to habitat of prey species:</u></p> <p>The impacts from nets, weights or anchors could cause limited damage to the habitat of the prey species of bottlenose dolphin Confidence in this assessment is medium (please see section 8).</p>
--	---

3. Feature description

<p>Feature Description:</p> <p>Bottlenose Dolphin</p>	<p>Bottlenose dolphin (<i>Tursiops truncatus</i>) is a large stocky cetacean that typically grows to around 2.5-2.7m in length and weighs between 200-275kg (Avant, 2008; Perrin <i>et al.</i>, 2009), although UK individuals are known to be larger than those found in the rest of the world, with individual maximum sizes of 4m recorded (JNCC, 2019a). Bottlenose dolphins have a worldwide distribution, primarily in tropical and temperate coastal and inshore regions (MarLIN, 2019). Populations around the UK are made up of two 'ecotypes': 'offshore' and 'coastal' (Wells and Scott, 1999). In the UK, group size for the coastal ecotype is commonly less than 20 animals, but groups of over 1000 animals have been recorded offshore (WDCS, 2002; Avant, 2008). Coastal bottlenose dolphin are of greater conservation concern in relation to fishing activities as they exhibit residency to inshore locations and are features of two Special Areas of Conservation (SACs) around the UK coast.</p> <p>The largest UK populations of coastal bottlenose dolphin are located in Cardigan Bay, Wales and the Moray Firth in north-east Scotland (JNCC, 2019a, 2019b), although smaller groups are found in Western Scotland and Southwest England. The Cardigan Bay Special Area of Conservation (SAC) was designated for coastal populations of bottlenose dolphin and this species is also a qualifying feature of Lleyn Peninsula and the Sarnau SAC (JNCC, 2019a, 2019b).</p> <p>Bottlenose dolphins have a broad diet, with a wide variety of benthic and pelagic fish species and invertebrates, including cephalopods. Their diet is thought to vary depending on availability of prey and they are considered an opportunistic and catholic feeder (Hernandez-Milian <i>et al.</i>, 2015).</p> <p>Females produce a single calf, every 2-3 years, in the summer after a gestation period of 12 months. The calf suckles for up to 18 months and stays close to the mother until it reaches four or five years of age (Avant, 2008; Perrin <i>et al.</i>, 2009).</p>
---	---

4. Gear description

<p>Gear Description:</p> <p>Fixed Trammel Nets</p>	<p>Fixed trammel nets are a type of gill net usually comprising three layers of monofilament or multifilament netting. Typically, an inner layer of fine small-mesh net (the lint) is sandwiched between two outer layers of large mesh nets (armouring) (Potter & Pawson, 1991). All three net layers are attached to a floated headline and weighted footrope, ensuring they hang vertically in the water, and the bottom of the net sits on or near the seabed (Potter & Pawson, 1991; FAO, 2019). The trammel net is fixed to the seabed at each end by conventional anchors or weights, thus preventing it moving in the tide, and nets are marked at one or both ends with buoys (Potter & Pawson, 1991; Seafish, 2019). Although trammel nets can be deployed in midwater or near the surface depending on design, buoyancy and target species (FAO, 2019), in the UK they are typically deployed on or just above the seabed (Seafish, 2019) and therefore the focus of this assessment is bottom-set fixed trammel nets.</p> <p>The inner small-mesh net of a trammel net is hung much slacker than the outer layers of netting, creating loose pockets of fine-mesh netting that enmesh or entangle a wider range of fish sizes and species, compared with standard gill nets (Potter & Pawson, 1991; Seafish, 2019). As with other types of gill net (gill, entangling and trammel), fish are typically (a) wedge-held, where the mesh catches around the body of the fish; (b) gill-held, when the mesh slips over the opercula; or (c) entangle-held, catching teeth, spines or other protrusions (Kalaycı & Yeşilçiçek, 2012).</p> <p>The use of net haulers is common on larger vessels hauling longer fleets of nets, often measuring several kilometres and/or nets set at greater depths (FAO, 2019). In small-scale inshore fisheries, as is common in Wales, individual trammel nets typically measuring a few hundred metres, and set in shallow or moderate depth water, could be hauled by hand or by net hauler.</p> <p>With all fixed net fisheries, a variety of international and national regulations and local factors determine the mesh size, length, and height of nets used, including areas fished and target species (NOAA, 2019). Relevant to Wales, EU and Welsh Government regulations determine mesh sizes, net lengths, soak times, and areas in which they can be deployed (Welsh Government, 2011a & 2011b; European Council, 2013).</p>
--	--

5. Assessment of impact pathways

<p>Assessment of impact pathway 1</p>	<p>1. Direct capture, damage, disturbance or harm to a designated species feature</p> <p>No studies were found that directly measured or estimated impacts of fixed trammel nets on bottlenose dolphin. Therefore, indirect evidence on the impacts from other fixed net fisheries on the direct capture, damage, disturbance or harm of bottlenose dolphin and similar species is considered.</p> <p>In Wales bottlenose dolphin was assessed to be in favourable condition during the 2018 Welsh indicative site level feature condition assessment for Cardigan Bay and Lleyn Peninsula and Sarnau SACs (NRW, 2018a; NRW, 2018b). However, Lohrengel <i>et al.</i> (2018) reported a significant decadal population decline for the wider Cardigan Bay area, a trend that has prompted further investigation into possible causes.</p> <p>The distribution of bottlenose dolphins overlaps with fixed net fishing activity predominantly by under 12m vessels within Welsh inshore waters (0-12NM), leading to the potential for bycatch interactions through entanglement by their teeth, beak, fins or tail (Bearzi <i>et al.</i>, 2008; Evans & Hintner, 2012; Baines & Evans, 2012; Zappes <i>et al.</i>, 2016).</p> <p>Globally, the magnitude of bottlenose dolphin bycatch is generally unknown or underreported (Reeves <i>et al.</i>, 2013). A report on dolphin by-catch across Europe by fixed net fisheries between 2006 and 2014 concluded, with low confidence, that by-catch was low and insufficient data was captured (Read <i>et al.</i>, 2017). Observational reports during the last decade (2011-2018) recorded a UK bycatch rate of one bottlenose dolphin per 6,292 hauls from predominantly offshore UK fixed gill net fisheries (Northridge <i>et al.</i>, 2011-2018). The UK Cetacean Strandings Investigation Programme (CSIP) have reported the cause of death of two bottlenose dolphins strandings in Wales between 2005 and 2015 was probably the result of trauma through bycatch with unidentified fishing gear (Deaville & Jepson, 2011; Deaville, 2012-16). However, it is not possible to identify the bycatch location from a stranded animal (ten Doeschate <i>et al.</i>, 2019).</p> <p>Bottlenose dolphin forage, locate fish, communicate and navigate using echolocation (Nowacek, 2005; Qunitana-Rizzo <i>et al.</i>, 2006) and are thought to be capable of detecting mono and multifilament bottom set gill nets from 25-55m in quiet conditions where there is zero to little background noise, and when approaching the net in a perpendicular direction (Kastelein <i>et al.</i>, 2000). This distance was thought to be lower when approaching from other angles, in noisier conditions, or where the dolphin might be distracted by movements of fish already caught in the net. This detection distance was thought sufficient to allow bottlenose dolphin time to react and take evasive action, when approaching a fixed net, therefore decreasing the likelihood of entrapment (Kastelein <i>et al.</i>, 2000).</p>
--	---

Mesh size, twine diameter, net height and water depth were identified as significant factors affecting cetacean bycatch rates in fixed nets (Palka and Rossman, 2001; Wiedenfeld *et al.*, 2015) and several of these factors were considered important for future research as potential bycatch mitigation measures (Wiedenfeld *et al.*, 2015; Northridge *et al.*, 2011-18). Palka and Rossman (2001) demonstrated United States bottlenose dolphin bycatch in fixed gillnet fisheries was 10-30 times greater for large mesh (≥ 7 inches, ≥ 178 mm) nets compared to the small (≤ 5 inches, ≤ 127 mm) net category and 4-9 times higher comparing large (≥ 7 inches, ≥ 178 mm) to medium (5-7 inches, 127-178mm) net sizes.

Kastelein *et al.* (2000) reported bottlenose dolphin's ability to detect different types of bottom set gill nets was not affected greatly by mesh size. They found that smaller mesh-nets with thinner twine but a greater density of knots and larger mesh sizes with fewer knots but thicker twine both provided strong return echolocation signals (Kastelein *et al.*, 2000). However, it should be noted that ability to detect nets was not linked to bycatch rates in this study. Fixed structures in the sea could act as barriers or deterrents, causing possible displacement or change in behaviour of bottlenose dolphin from an otherwise suitable habitat (Shane *et al.*, 1986; Markowitz, 2004). The operation of fixed nets is usually temporary, however, depending on the location and amount of fixed netting the activity could potentially cause barrier or deterrent effects.

In order to deter bottlenose dolphin, Regulation (EU) 2019/1241 (European Union, 2019) requires vessels of ≥ 12 m in certain areas (excluding the Irish Sea - ICES area 7a), to use active acoustic deterrent devices emitting high-frequency pulsed sounds (pingers) on specified fishing gears. The effectiveness of pingers to deter bottlenose dolphin is in doubt (Cox *et al.*, 2003; Gazo *et al.*, 2008; Waples *et al.*, 2013). A small-scale study, observing 69 bottlenose dolphin concluded caution in the use of pingers, stating they would be unlikely to reduce bottlenose dolphin bycatch in gill net fisheries (Cox *et al.*, 2003). Additionally, Cox *et al.* (2003) stated that bottlenose dolphin, in their study, displayed behaviour supporting the 'dinner bell effect' originally described by Mate and Harvey (1987), where cetaceans or seals become sensitised to pingers over time and learn to associate the pinger sound with a source of food.

Activities that produce underwater noise have the potential to disturb bottlenose dolphin. It is difficult to separate the behavioural changes in cetaceans due to the additional noise of fishing vessels versus only background noise levels (Rako-Gospic and Picciulin, 2016). Studies show that small vessels produce sounds which overlap in frequencies with bottlenose dolphin vocalisation involved in social interactions (Rako-Gospic and Picciulin, 2016). This type of 'auditory masking' may therefore influence the behaviour of groups of bottlenose dolphin, including causing 'seasonal displacement' when boat traffic is high (Rako *et al.*, 2013; Peng *et al.*, 2015).

Commercial fishing contributes to ambient noise including low frequency sound emitted from engines and gear winching and hauling, and high frequency sound from the use of sonar and fish finding equipment (Evans and Hintner, 2012). Bottlenose dolphin have been observed responding to boats in different ways, depending on the type of vessel and noise (Rako-Gospic and Picciulin, 2016) and how the vessel interacts with the dolphin

	<p>(Lusseau, 2006). In Cardigan Bay, bottlenose dolphin were observed responding to approaching boats at a distance of 150-300m by making longer dives and moving away from the source of the sound (Pesante <i>et al.</i>, 2008b). In contrast, bottlenose dolphin can be attracted to boats and display bow riding behaviour (Williams <i>et al.</i>, 1992), whereas foraging individuals are likely to be aware of, but will ignore vessels (Richardson <i>et al.</i>, 1995). However, these types of interactions occur mostly with moving vessels and are not as relevant to the static gear itself.</p> <p>Depending on the fishery, the operation of the gear and the intensity of the activity it is possible that the impacts from fixed trammel nets or noise pollution associated with fishing vessels could lead to bottlenose dolphin bycatch, displacement or disturbance.</p>
<p>Assessment of impact pathway 2</p>	<p>2. Damage to the habitat of designated species features (including through direct physical impact, pollution, changes in thermal regime, hydrodynamics, light etc.)</p> <p>No studies were found that directly measured or estimated the impacts of fixed trammel nets on the habitat utilised by bottlenose dolphin. Therefore, indirect evidence on the impacts from other fixed nets on the habitats utilised by bottlenose dolphin is considered.</p> <p>Bottlenose dolphin are a highly mobile species exhibiting a full spectrum of movements, including seasonal migrations, year-round home ranges, periodic residency, and a combination of occasional long-range movements and repeated local residency (Reid <i>et al.</i>, 2003; Feingold and Evans, 2014b; Lohrengel <i>et al.</i>, 2018). They use both inshore and offshore areas of Wales (Pesante <i>et al.</i>, 2008a, 2008b; Feingold and Evans, 2014a and 2014b; Lohrengel <i>et al.</i>, 2018) with a high frequency of sightings along the coast from Aberaeron to Cardigan and around Fishguard suggesting that these coastal areas may be of particular significance (Baines and Evans, 2012).</p> <p>Defining the specific habitat requirements for cetaceans is difficult due to their wide-ranging and highly mobile nature and their distribution is likely driven by the corresponding distribution and availability of their various prey species, and other unknown factors (NRW and JNCC, 2017). Important habitat considerations for bottlenose dolphin include the seabed and water column habitats of sufficient quality for feeding, breeding and calving, socialising and resting (Lopes, 2017). The impact of changes to the physical habitat of bottlenose dolphin from human activity on population size has not been observed (Shane <i>et al.</i>, 1986).</p> <p>Prey availability is thought to be the main driver for bottlenose dolphin distribution as they tend to be found in areas of high marine productivity (Lopes, 2017). In UK coastal waters, bottlenose dolphin appears to favour habitat with uneven topography and strong tidal currents. Acoustic monitoring has demonstrated the importance of sandbanks, coarse sediments and reefs for foraging (Pesante <i>et al.</i>, 2008a, 2008b; Feingold and Evans,</p>

	<p>2014a; Lopes, 2017). The interaction between fixed entangling nets and the benthic habitats of bottlenose dolphin prey is considered in impact pathway 4.</p> <p>Bottlenose dolphin activities, other than benthic feeding, tend to occur within the water column and are not known to be dependent on the seabed habitat. The impacts from nets, weights or anchors are not likely to affect the integrity of the water column habitats utilised by bottlenose dolphin.</p>
<p>Assessment of impact pathway 3</p>	<p>3. Removal of prey species of a designated species feature</p> <p>No studies were found that directly measured or estimated the impacts of fixed trammel nets removing the prey species of bottlenose dolphin. Therefore, indirect evidence from other fixed net fisheries catching the prey of bottlenose dolphin can be considered.</p> <p>Evidence indicates bottlenose dolphins target some of the same species that fixed trammel net fisheries target (Reid <i>et al.</i>, 2003; Hernandez-Milian <i>et al.</i>, 2015; Seafish, 2011; Seafish, 2019; Walmsley & Pawson, 2007; Tregenza <i>et al.</i>, 1997). Competition is likely to occur between commercial fishing activities and foraging bottlenose dolphin (Hernandez-Milian <i>et al.</i>, 2015). However, bottlenose dolphin, in common with other small marine mammal species that switch diets and feed in ecosystems where the choice of prey is varied, are less likely to be dramatically affected by fishing impacts on their prey species (Reid <i>et al.</i>, 2003; Hernandez-Milian <i>et al.</i>, 2015; Giménez <i>et al.</i>, 2017; Hutchinson, 1996; Jennings <i>et al.</i>, 2001). Overlapping prey target species between the bottlenose dolphin and fixed trammel net fisheries include sole, plaice, dab and flounder within the inner section of the net (mesh sizes of 100-120mm) (Walmsley & Pawson, 2007) and hake, pollock, rays, turbot, monkfish and brill within the larger outer nets (mesh sizes of 200-300mm) (Tregenza <i>et al.</i>, 1997).</p> <p>The Maximum Sustainable Yield (MSY) for a fish stock is the maximum level at which a fish stock can be routinely exploited without long-term depletion. In the pursuit of MSY for fish stocks, the International Council for the Exploration of the Seas (ICES) incorporates both fishing and natural fish mortality in their multi-species stock assessment models and Total Allowable Catch (TAC) advice for North Sea cod, haddock, herring, whiting, sprat, Norway pout and sandeel (Walmsley, 2018). Natural mortality is defined as “all sources of mortality of a fish stock outside of that caused by fishing” (Walmsley, 2018). Specifically, this includes predation by other fish, birds and marine mammals, and mortality from biotic and abiotic factors such as temperature, disease and from other anthropogenic activities, excluding fishing (Walmsley, 2018).</p> <p>ICES have recently developed a multi-species model for the North Sea, including cod, haddock, herring, whiting, sprat, Norway pout and sandeel (Walmsley, 2018). A similar multi-species stock assessment model is currently being developed for the Irish Sea (Walmsley, 2018), and once complete, should incorporate predator/prey species interactions, such as those from foraging bottlenose dolphin. Once this improved estimate of natural</p>

	<p>mortality is incorporated into stock assessments and TAC advice in the Irish Sea, the impact of removing bottlenose dolphin prey species by fixed entangling nets will be of lesser importance to bottlenose dolphin populations. However, importantly, it should be noted that non-commercial fish species forming part of the diet of bottlenose dolphin are not subject to the same stock assessments.</p> <p>Depending on the intensity of fishing activity, it is possible that the removal of prey species by fixed trammel nets could affect bottlenose dolphins. However, evidence suggests that bottlenose dolphin will readily switch prey, but it is not known if dependency on alternative prey availability and quality is detrimental at the population level in the long term.</p>
<p>Assessment of impact pathway 4</p>	<p>4. Damage to habitat of prey species</p> <p>No studies were found that directly measured or estimated the impacts of fixed trammel nets on the habitats of prey species of bottlenose dolphins. Therefore, indirect evidence on the impacts from other fixed net fisheries on the habitats utilised by bottlenose dolphin prey species is considered.</p> <p>Prey species of bottlenose dolphin include (but are not limited to) cephalopods and other shellfish and demersal and pelagic fish including, bass, blue whiting, cod, eels, flatfish, haddock, hake, mullet, octopus, pollock, saithe, salmon, sandeels, silver pout, snipefish, sprat, trout and whiting (Reid <i>et al.</i>, 2003; Hernandez-Milian <i>et al.</i>, 2015). The habitat of these prey species varies but can be broadly characterised as pelagic and benthic habitats including sediments such as sands and gravels, seagrass beds and reefs.</p> <p>In UK coastal waters, bottlenose dolphin appears to favour habitat with uneven topography and strong tidal currents, where their prey species occur. Acoustic monitoring has demonstrated the importance of sedimentary habitats and reefs for foraging (Pesante <i>et al.</i>, 2008a, 2008b; Feingold and Evans, 2014a; Lopes, 2017). Sedimentary habitats, located in bays and estuaries include sandbanks, and in more sheltered environments, seagrass beds. These are considered important habitat and nursery areas for various demersal and pelagic fish species (Bertelli and Unsworth, 2014), many of which are prey species for bottlenose dolphin. Anchors and weights, distributed along the foot rope of fixed entangling nets, have the potential to penetrate finer sediments including sands and gravels, and nets set in higher-energy environments may cause greater abrasion to the seabed due to the increased tidal forces acting on the nets. In sand and gravel habitats the mobile and dynamic nature of the seabed (Hinz <i>et al.</i>, 2010a and 2010b; JNCC, 2017) combined with the relatively small footprint of fixed net anchors, the relatively short soak times of the nets measured in hours (Northridge <i>et al.</i>, 2017) all suggest that any seabed disturbances from anchors and weights are likely to recover over short periods of time, such as weeks (Hinz <i>et al.</i>, 2010a and 2010b).</p> <p>Reef habitats are potentially at risk of abrasion or crushing by fixed net anchors or weights, nets can also become entangled on seabed structures causing fragmentation, tearing or abrasion of the habitat, leading to deterioration and the removal of long-lived fragile and emergent epifauna (Brown and Macfadyen, 2007).</p>

	<p>However, most fishers with nets tend to avoid reef habitats to prevent losing or damaging their nets, and so the risk to these habitats may be lower than anticipated.</p> <p>Factors affecting the integrity of pelagic fish habitats e.g. water quality are not likely to be affected by fishing with fixed trammel nets. These factors are not considered further in this assessment.</p> <p>Depending on the footprint and the intensity of the activity it is possible the impacts from nets, weights or anchors could cause limited damage to the habitat of the prey species of bottlenose dolphin.</p> <p>However, these are large scale habitat features and there is no evidence to suggest that the impacts from fixed net fisheries on the habitats of bottlenose dolphin prey species would affect the bottlenose dolphin at a population level.</p>
--	--

6. SACs designated for bottlenose dolphin

Bottlenose dolphin are listed as protected species in two Welsh SACs, but due to their mobile nature, impacts from activities must be considered throughout their wider management unit.

<p>Cardigan Bay SAC</p>	<p>Bottlenose dolphin are often sighted within Cardigan Bay SAC off headlands and in more sheltered areas near New Quay, Ynys Lochtyn, Aberporth, Mwnt, and the Teifi Estuary. It should be noted that the coast between New Quay and Cemaes Head has been the area of greatest observer effort over the years (Lohrengel <i>et al.</i>, 2018; NRW, 2018c).</p> <p>Recent surveys show that the numbers of bottlenose dolphin are greatest from July and October with fewer seen between November and April, although some animals are present near shore in every month of the year. They are most commonly seen within 10 miles of the coast, from April to October and most concentrated within 2 miles near headlands, estuaries and in embayments (NRW, 2018c).</p> <p>Bottlenose dolphin was assessed to be in favourable condition during the latest 2018 SAC feature condition assessment for the Cardigan Bay SAC (NRW, 2018a, c). However, Lohrengel <i>et al.</i> (2018) also reported a significant decadal population decline for the wider Cardigan Bay area, a trend that has prompted further investigation into possible causes.</p> <p>Bottlenose dolphin using the SAC are part of the wider Irish Sea Management Unit (IAMMWG, 2015; Lohrengel <i>et al.</i>, 2018).</p>
<p>Lleyn Peninsula and the Sarnau SAC</p>	<p>The Lleyn Peninsula and Sarnau SAC supports the same population of bottlenose dolphin found in Cardigan Bay SAC. Bottlenose dolphin using the SAC are part of the wider Irish Sea Management Unit (IAMMWG, 2015; Lohrengel <i>et al.</i>, 2018).</p> <p>Bottlenose dolphin was assessed to be in favourable condition during the latest 2018 SAC feature condition assessment for the Lleyn Peninsula and Sarnau SAC (NRW, 2018b, d). However, Lohrengel <i>et al.</i> (2018) also reported a significant decadal population decline for the wider Cardigan Bay area, a trend that has prompted further investigation into possible causes.</p> <p>Bottlenose dolphin using the SAC are part of the wider Irish Sea Management Unit (IAMMWG, 2015; Lohrengel <i>et al.</i>, 2018).</p>

7. Evidence Gaps

- Direct studies to measure the impacts from fixed trammel nets on bottlenose dolphins
- Studies to measure noise pollution of Welsh fishing fleet on bottlenose dolphin
- Studies to measure behaviour change of bottlenose dolphin towards pingers
- Monitoring of bottlenose dolphin population status and structure

8. Confidence assessment

The confidence score is the sum of scores from three evidence components: quality, applicability and agreement. These are qualitatively assessed as high, medium or low using the most appropriate statements in the table below, and these are numerically represented as scores of 3, 2, or 1 respectively.

A total confidence score of 3 – 4 represents low confidence, 5 – 7 shows medium confidence and 8 – 9 demonstrates high confidence in the evidence used in the assessment.

This assessment scores 7, representing medium confidence in the evidence.

	Evidence quality	Evidence applicability	Evidence agreement
High	Based on more than 3 recent and relevant peer reviewed papers or grey literature from established agencies.	Based on the fishing gear acting on the feature in the UK.	Strong agreement between multiple (>3) evidence sources. Score 3.
Medium	Based on either relevant but older peer reviewed papers or grey literature from less established agencies; or based on only 2-3 recent and relevant peer reviewed evidence sources. Score 2.	Based on similar fishing gears, or other activities with a similar impact, acting on the feature in the UK. Score 2.	Some disagreement but majority of evidence agrees. Or fewer than 3 evidence sources used.
Low	Based on either less relevant or older grey literature from less established agencies; or based on only 1 recent and relevant peer reviewed evidence source.	Based on dissimilar fishing gears acting upon the feature in other areas.	Little agreement between evidence.

N.B. When evidence is indirect the evidence quality and applicability will be capped to medium, to ensure that direct evidence gaps are captured in this approach.

9. References

- Avant, P. (2008). *Tursiops truncatus* Bottle-nosed dolphin. In Tyler-Walters H. and Hiscock K. (eds) Marine Life Information Network: Biology and Sensitivity Key Information Reviews, [on-line]. Plymouth: Marine Biological Association of the United Kingdom.
<http://www.marlin.ac.uk/species/detail/1674>.
- Baines, M.E. and Evans, P.G.H. (2012). Atlas of the Marine Mammals of Wales. 2nd Edition. Marine Monitoring Report No. 68. Countryside Council for Wales, Bangor.
- Bearzi, G., Fortuna, C.M. and Reeves, R.R. (2008). Ecology and conservation of common bottlenose dolphins *Tursiops truncatus* in the Mediterranean Sea. Mammal Rev. 2008, Volume 39, No. 2, 92-123.
- Bertelli, C.M. and Unsworth, R.K.F. (2014). Protecting the hand that feeds us: Seagrass (*Zostera marina*) serves as commercial juvenile fish habitat. Marine Pollution Bulletin, Vol. 83, Issue 2: 425-429.
- Brown, J. and Macfadyen, G. (2007). Ghost fishing in European waters: Impacts and management responses. Marine Policy, Elsevier, vol. 31(4), pages 488-504.
- Cox, T.M., Read, A.J., Swanner, D., Urian, K. and Waples, D. (2003). Behavioural responses of bottlenose dolphins, *Tursiops truncatus*, to gill nets and acoustic alarms. Biological Conservation, 115, 203–212.
- Deville, R. and Jepson, P.D. (2011). Final report for the period 1st January 2005 - 31st December 2010. *UK Cetacean Strandings Investigation Programme (CSIP)*.
- Deville, R. (2012). Final report for the period 1st January - 31st December 2011. UK Cetacean Strandings Investigation Programme (CSIP).
[http://randd.defra.gov.uk/Document.aspx?Document=11149_FINALUKCSIPAnnualReport2011\(2\).pdf](http://randd.defra.gov.uk/Document.aspx?Document=11149_FINALUKCSIPAnnualReport2011(2).pdf).
- Deville, R. (2013). Final report for the period 1st January - 31st December 2012. UK Cetacean Strandings Investigation Programme (CSIP).
http://randd.defra.gov.uk/Document.aspx?Document=11840_FINALUKCSIPAnnualReport2012.pdf.

Deaville, R. (2014). Final report for the period 1st January - 31st December 2013. UK Cetacean Strandings Investigation Programme (CSIP). http://randd.defra.gov.uk/Document.aspx?Document=12306_UKCSIPAnnualReport2013_Final.pdf.

Deaville, R. (2015). Final report for the period 1st January - 31st December 2014. UK Cetacean Strandings Investigation Programme (CSIP). http://randd.defra.gov.uk/Document.aspx?Document=12562_Final_UK_CSIP_Annual_Report_2014.pdf.

Deaville, R. (2016). Final report for the period 1st January - 31st December 2015. UK Cetacean Strandings Investigation Programme (CSIP). http://randd.defra.gov.uk/Document.aspx?Document=14001_FINALUKCSIPAnnualReport2015.pdf.

European Council. (2013). Council Regulation (EC) No 227/2013 for the conservation of fishery resources through technical measures for the protection of juveniles of marine organisms. <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:31998R0850>.

European Union. (2019). Regulation (EU) 2019/1241 of the European Parliament and of the Council of 20 June 2019 on the conservation of fisheries resources and the protection of marine ecosystems through technical measures, amending Council Regulations (EC) No 1967/2006, (EC) No 1224/2009 and Regulations (EU) No 1380/2013, (EU) 2016/1139, (EU) 2018/973, (EU) 2019/472 and (EU) 2019/1022 of the European Parliament and of the Council, and repealing Council Regulations (EC) No 894/97, (EC) No 850/98, (EC) No 2549/2000, (EC) No 254/2002, (EC) No 812/2004 and (EC) No 2187/2005. <https://eur-lex.europa.eu/legal-content/en/TXT/?uri=CELEX:32019R1241>.

Evans, P.G.H. and Hintner, K. (2012). A review of the direct and indirect impacts of fishing activities on marine mammals in Welsh waters. CCW Policy Research Report No. 12/5: 1-172.

FAO. (2019). Fishing Gear types. Gillnets and entangling nets. Fisheries and Aquaculture Department Technology Fact Sheets. [Accessed 12th December 2019]. <http://www.fao.org/fishery/geartype/107/en>.

Feingold, D. and Evans P.G.H. (2014a). Bottlenose Dolphin and Harbour Porpoise Monitoring in Cardigan Bay and Pen Llŷn a'r Sarnau Special Areas of Conservation 2011 - 2013. NRW Evidence Report Series Report No: 4, 120 pp, Natural Resources Wales.

Feingold, D. and Evans, P.G.H. (2014b). Connectivity of Bottlenose Dolphins in Welsh Waters: North Wales Photo-Monitoring Report. Sea Watch Foundation.

Gazo, M., Gonzalvo, J. and Aguilar, A. (2008). Pingers as deterrents of bottlenose dolphins interacting with trammel nets. Fisheries Research, 92, 70–75.

Giménez, J., Marcalo, A., Ramirez, P.V., Gauffier, P., Esteban, R., Nicolau, L., Gonzalez-Ortegon, E., Baldo, F., Vilas, C., Vingada, J., Forero, M.G. and de Stephanis, R. (2017). Diet of Bottlenose dolphins (*Tursiops truncatus*) from the Gulf of Cadiz: Insights from stomach content and stable isotope analyses. PLoS ONE 12(9): e0184673. <https://doi.org/10.1371/journal.pone.0184673>.

He, P. (2006). Gill nets: gear design, fishing performance and conservation challenges. Marine Technology Society Journal 40(3):12-19.

Hernandez-Milian, G., Berrow, S., Santos, M., Reid, D. and Rogan, E. (2015). Insights into the trophic ecology of bottlenose dolphins (*Tursiops truncatus*) in Irish waters. Aquatic Mammals, 41(2): 226-239.

Hinz, H., Scriberras, M., Benell, J.D. and Kaiser, M.J. (2010a). Assessment of offshore habitats in the Cardigan Bay SAC. Fisheries and Conservation report, Bangor University, p.22.

Hinz, H., Scriberras, M., Murray, L.G., Benell, J.D. and Kaiser, M.J. (2010b). Assessment of offshore habitats in the Cardigan Bay SAC (June 2010 survey). Fisheries and Conservation report, (14), p.30.

Hutchinson, J.D. (1996). Fisheries interactions: the harbour porpoise – a review. In: The conservation of whales and dolphins: Science and Practice. (eds M.P. Simmonds and J.D. Hutchinson), pp.128-165. John Wiley, Chichester.

IAMMWG. (2015). Management Units for cetaceans in UK waters (January 2015). JNCC Report No. 547, JNCC Peterborough.

Jennings, S., Kaiser, M.J. and Reynolds, J.D. (2001). *Marine Fisheries Ecology*. Blackwell Science. ISBN 0-632-05098-5.

JNCC. (2017). Annex 1 Sandbanks in offshore waters. [Viewed 21/08/2017]. <http://jncc.defra.gov.uk/page-1452>.

JNCC. (2019a). Annex II Species accounts: Bottlenose dolphin *Tursiops truncatus*. [Viewed 13 February 2019]. <http://jncc.defra.gov.uk/ProtectedSites/SACselection/species.asp?FeatureIntCode=s1349>.

JNCC. (2019b). European Community Directive on the Conservation of Natural Habitats and of Wild Fauna and Flora (92/43/EEC) Fourth Report by the United Kingdom under Article 17 on the implementation of the Directive from January 2013 to December 2018 Conservation status assessment for Species: S1349 – Bottlenose Dolphin (*Tursiops truncatus*). <https://jncc.gov.uk/jncc-assets/Art17/S1349-UK-Habitats-Directive-Art17-2019.pdf>.

- Kalaycı, F. and Yeşilçiçek, T. (2012). Investigation of the selectivity of trammel nets used in red mullet (*Mullus barbatus*) fishery in the eastern Black Sea, Turkey. *Turkish Journal of Fisheries and Aquatic Sciences*, 12(4).
- Kastelein, R.A., Au, W.W.L. and de Haan, D. (2000). Detection distances of bottom-set gill nets by harbour porpoises (*Phocoena phocoena*) and bottlenose dolphins (*Tursiops truncatus*). *Marine Environmental Research* 49:359-375.
- Lohrengel, K., Evans, P.G.H., Lindenbaum, C.P., Morris, C.W. and Stringell, T.B. (2018). Bottlenose Dolphin Monitoring in Cardigan Bay 2014 - 2016. NRW Evidence Report No: 191, 162pp, Natural Resources Wales, Bangor.
- Lopes, K.F.C.F. (2017). Habitat Preference of Bottlenose Dolphin (*Tursiops truncatus*) in Cardigan Bay. MSc Thesis. Bangor University. http://www.seawatchfoundation.org.uk/wp-content/uploads/2017/11/Katia-Lopes_MSc-thesis.pdf.
- Lusseau, D. (2006). The short-term behavioural reactions of bottlenose dolphins to interactions with boats in Doubtful Sound, New Zealand. *Marine Mammal Science*, 22(4), pp.802-818.
- Markowitz, T.M. (2004). *Social organization of the New Zealand dusky dolphin* (Doctoral dissertation, Texas AandM University).
- MarLIN. (2019). Bottle-nosed dolphin (*Tursiops truncatus*). Marine Life Information Network (MarLIN). [Viewed 13 February 2019]. <https://www.marlin.ac.uk/species/detail/1674>.
- Mate, B.R. and Harvey, J.T. (1987). Acoustical deterrents in marine mammal conflicts with fisheries. Oregon Sea Grant Report ORESU-W-86-001. 116 pp.
- NOAA. (2019). Fishing Gear: Gillnets. NOAA Fisheries. [Accessed 05th March 2019]. <https://www.fisheries.noaa.gov/national/bycatch/fishing-gear-gillnets>.
- Northridge, S., Kingston, A. and Thomas, L. (2011). Annual report on the implementation of Council Regulation (EC) No 812/2004 during 2010.
- Northridge, S., Kingston, A. and Thomas, L. (2012). Annual report on the implementation of Council Regulation (EC) No 812/2004 during 2011.
- Northridge, S., Kingston, A. and Thomas, L. (2013). Annual report on the implementation of Council Regulation (EC) No 812/2004 during 2012.

- Northridge, S., Kingston, A. and Thomas, L. (2014). Annual report on the implementation of Council Regulation (EC) No 812/2004 during 2013.
- Northridge, S., Kingston, A. and Thomas, L. (2015). Annual report on the implementation of Council Regulation (EC) No 812/2004 during 2014.
- Northridge, S., Kingston, A. and Thomas, L. (2016). Annual report on the implementation of Council Regulation (EC) No 812/2004 during 2015.
- Northridge, S., Kingston, A. and Thomas, L. (2017). Annual report on the implementation of Council Regulation (EC) No 812/2004 during 2016.
- Northridge, S., Kingston, A. and Thomas, L. (2018). Annual report on the implementation of Council Regulation (EC) No 812/2004 during 2017.
- Nowacek, D.P. (2005). Acoustic ecology of foraging bottlenose dolphins (*Tursiops truncatus*), habitat-specific sound use of three sound types. *Marine Mammal Science*. 21, 587-602.
- NRW and JNCC. (2017). SAC Selection Assessment Document: West Wales Marine / Gorllewin Cymru Forol. January, 2017. Natural Resources Wales and Joint Nature Conservation Committee, UK.
- NRW. (2018a). Cardigan Bay / Bae Ceredigion Special Area of Conservation: Indicative site level feature condition assessments 2018. NRW Evidence Report Series, Report No: 226, 39pp, NRW, Bangor.
- NRW. (2018b). Pen Llŷn a'r Sarnau / Lleyn Peninsula and the Sarnau Special Area of Conservation: Indicative site level feature condition assessments 2018. NRW Evidence Report Series, Report No: 234, 58pp, NRW, Bangor.
- NRW. (2018c). Cardigan Bay/ Bae Ceredigion Special Area of Conservation. Advice provided by Natural Resources Wales in fulfilment of Regulation 37 of the Conservation of Habitats and Species Regulations 2017. Natural Resources Wales, Bangor pp 87.
- NRW. (2018d). Pen Llŷn a'r Sarnau/Lleyn Peninsula and the Sarnau Special Area of Conservation Advice provided by Natural Resources Wales in fulfilment of Regulation 37 of the Conservation of Habitats and Species Regulations 2017. Natural Resources Wales, Bangor pp 143.
- Palka, D.L. and Rossman, M.C. (2001). Bycatch estimates of coastal bottlenose dolphin (*Tursiops truncatus*) in U.S. Mid-Atlantic gillnet fisheries for 1996 to 2000. Northeast Fisheries Science Centre NEFSC. Reference Document 01-15; 77p.

Peng, C., Zhao, X., and Liu, G. (2015). Noise in the sea and its impacts on marine organisms. *International journal of environmental research and public health*, 12(10), 12304-12323.

Perrin, W.F., Würsig, B. and Thewissen, J.G.M. eds. (2009). *Encyclopaedia of marine mammals*. Academic Press.

Pesante, G., Evans, P.G.H., Anderwald, P., Powell, D. and McMath, M. (2008a). Connectivity of bottlenose dolphins in Wales: North Wales photo-monitoring. CCW Marine Monitoring Report No. 62. Countryside Council for Wales, Bangor.

Pesante, G., Evans, P.G.H., Baines, M.E. and McMath, M. (2008b). Abundance and Life History Parameters of Bottlenose Dolphin in Cardigan Bay: Monitoring 2005-2007. CCW Marine Monitoring Report No. 61. Countryside Council for Wales, Bangor.

Potter, E.C.E., and Pawson, M.G. (1991). *Gill netting*. Ministry of Agriculture, Fisheries and Food, Directorate of Fisheries Research.

Qunitana-Rizzo, E., Mann, D.A. and Wells, R.S. (2006). Estimated communication range of social sounds used by bottlenose dolphins (*Tursiops truncatus*). *J. Acoust. Soc. Am.* 120, 1671-1683.

Rako, N., Fortuna, C.M., Holcer, D., Mackelworth, P., Nimak-Wood, M., Pleslic, G., Sebastianutto, L., Vilibic, I., Wiemann, A. and Picciulin, M. (2013). Leisure boating noise as a trigger for the displacement of the bottlenose dolphins of the Cres–Lošinj archipelago (northern Adriatic Sea, Croatia). *Marine Pollution Bulletin* 68 (2013) 77–84.

Rako-Gospić, N. and Picciulin, M. (2016). Changes in whistle structure of resident bottlenose dolphins in relation to underwater noise and boat traffic. *Marine Pollution Bulletin* 105 (2016) 193–198.

Read, F.L., Evans, P.G.H. and Dolman, S.J. (2017). Cetacean Bycatch Monitoring and Mitigation under EC Regulation 812/2004 in the Northeast Atlantic, North Sea and Baltic Sea from 2006 to 2014. A WDC Report. 68 pp.

Reeves, R.R., McClellan, K. and Werner, T.B. (2013). Marine mammal bycatch in gillnet and other entangling net fisheries, 1990 to 2011. *Endangered Species Research* Vol. 20: 71–97.

Reid, J.B., Evans, P.G.H. and Northridge, S.P. (2003). Atlas of Cetacean distribution in north-west European waters. Joint Nature Conservation Committee, Peterborough.

Richardson, W.J., Greene, C.R.J., Malme, C.I. and Thomson, D.H. (1995). *Marine Mammals and Noise*. San Diego, Academic Press.

Seafish. (2011). Gear Technology Note – Static Gear, February 2011.

http://www.seafish.org/media/Publications/SeafishGuidanceNote_StaticGear_201102.pdf.

Seafish. (2019). Fishing Gear Database: Tangle Nets. [Accessed 12th December 2019]. <https://www.seafish.org/gear/gear/profile/tangle-nets>.

Shane, S.H., Wells, R.S., and Würsig, B. (1986). Ecology, behavior and social organization of the bottlenose dolphin: a review. *Marine mammal science*, 2(1), 34-63.

ten Doeschate, M.T.I., Brownlow A.C., Morris C.W. and Stringell T.B. (2019). UK West Coast Harbour Porpoise (*Phocoena phocoena*) Strandings. NRW Evidence Report Series. Report No: 358. 66pp. Natural Resources Wales, Bangor.

Tregenza, N.J.C., Berrow, S.D., Hammond, P.S. and Leaper, R. (1997). Harbour porpoise (*Phocoena* L.) by-catch in set gill nets in the Celtic Sea. – ICES Journal of Marine Science, 54: 896–904.

Walmsley, S.A. and Pawson, M.G. (2007). The coastal fisheries of England and Wales, Part V: a review of their status 2005-6. Science Series Technical Reports, Cefas Lowestoft, 140: 83pp.

Walmsley, S.F. (2018). Marine Fish Mortality Considerations as part of Maximum Sustainable Yield Calculations NRW Report No: 267, 37pp, Natural Resources Wales, Bangor.

Waples, D.M., Thorne, L.H., Hodge, L.E.W., Burke, E.K., Urian, K.W. and Read, A.J. (2013). A field test of acoustic deterrent devices used to reduce interactions between bottlenose dolphins and a coastal gill net fishery. *Biological Conservation*, Volume 157, Pages 163-171.

WDCS. (2002). [Cited 12 July 2017]. <http://www.wdcs.org/dan/publishing.nsf/allweb/4BE4E4DD1F6DF314802568F8004F381E>.

Wells, R.S. and Scott, M.D. (1999). Bottlenose Dolphin. In: *Handbook of Marine Mammals*. Volume 6: The Second Book of Dolphins and Porpoises. Pp 137-182. Academic Press, San Diego. 486pp.

Welsh Government. (2011a). Inshore Fishery Legislation: South Wales 0-6 nautical miles. Text of the saved Byelaws of the former South Wales Sea Fisheries Committee.

Welsh Government. (2011b). Inshore Fishery Legislation: North Wales 0-6 nautical miles. Text of the saved Byelaws of the former North Western and North Wales Sea Fisheries Committee.

Wiedenfeld, D.A., Crawford, R. and Pott, C.M. (2015). Results of a Workshop on Reduction of Bycatch of Seabirds, Sea Turtles, and Sea Mammals in Gill nets, 21-23 January 2015. American Bird Conservancy and Bird Life International.

Williams, T.M., Friedl, W.A., Fong, M.L., Yamada, R.M., Sedivy, P. and Haun, J.E. (1992). Travel at low energetic cost by swimming and wave-riding bottlenose dolphins. *Nature*, 355(6363), pp.821-823.

Zappes, C.A. Simoes-Lopes, P.C. Andriolo, A. and Madeira Di Benedetto, A.P. (2016). Traditional knowledge identifies causes of bycatch on bottlenose dolphins (*Tursiops truncatus* Montagu 1821): an ethnobiological approach. *Ocean and Coastal Management*, Volume 120, pp. 160-169.